

Advanced Stage at Diagnosis and Worse Clinicopathologic Features in Young Women with Breast Cancer in Brazil: A Subanalysis of the AMAZONA III Study (GBECAM 0115)

Maria Alice Franzoi, MD^{1,2,3}; Daniela D. Rosa, MD, PhD^{3,4,5}; Facundo Zaffaroni, MA³; Gustavo Werutsky, MD^{3,4}; Sérgio Simon, MD, PhD^{3,4}; José Bines, MD, PhD^{3,4,6}; Carlos Barrios, MD^{3,4}; Eduardo Cronemberger, MSc⁷; Geraldo Silva Queiroz, MD⁸; Vladimir Cordeiro de Lima, MD, PhD⁹; Ruffo Freitas Júnior, MD, PhD¹⁰; José Couto, PhD¹¹; Karla Emerenciano, MD¹²; Heloísa Resende, PhD¹³; Susanne Crocama, PhD¹⁴; Tomás Reinert, MD¹⁵; Brigitte Van Eyli, MD¹⁶; Yeni Nerón, MD¹⁷; Vanessa Dybal, MD¹⁸; Nicolas Lazaretti, MSc¹⁹; Rita de Cassia Costamillan, MD²⁰; Diocésio Alves Pinto de Andrade, MD²¹; Clarissa Mathias, MD, PhD²²; Giovana Zerwes Vacaro, MD²³; Giuliano Borges, MD²⁴; Alessandra Morelle, PhD⁵; Carlos Alberto Sampaio Filho, MD¹⁸; Max Mano, MD, PhD²⁵; and Pedro E. R. Liedke, MD, MSc^{1,3,4}

PURPOSE Breast cancer (BC) in young women is uncommon and tends to present with more aggressive characteristics. To better understand and characterize this scenario in Brazil through real-world data, we performed a subanalysis of AMAZONA III study (ClinicalTrials.gov identifier: NCT02663973).

METHODS The AMAZONA III study (GBECAM 0115) is a prospective registry that included 2,950 women newly diagnosed with invasive BC in Brazil from January 2016 until March 2018 at 22 sites. Valid data were obtained from 2,888 patients regarding age at diagnosis and complete baseline information. To compare epidemiologic and clinicopathological features at the time of diagnosis, patients with BC were divided into two groups according to age: ≤ 40 years and > 40 years. Quantitative variables were described as means, and categorical variables were described as frequencies and percentages and compared using the Pearson's χ^2 test.

RESULTS Of 2,888 women diagnosed with BC, 486 (17%) were ≤ 40 years old. Young women had higher educational level, most were employed and a significant number were married ($P < .001$ for all associations). Younger patients were more symptomatic at BC diagnosis ($P < .001$), and they also presented more frequently with stage III, T3/T4, grade 3 tumors, HER-2–positive, luminal B, and triple-negative subtypes.

CONCLUSION Brazilian women younger than age 40 years have unfavorable clinicopathological features of BC at diagnosis, with more aggressive subtypes and advanced stage when compared with older women. These differences are not explained by socioeconomic or ethnic imbalances. The causes of a higher prevalence of BC among young women in Brazil deserve additional investigation.

J Global Oncol. © 2019 by American Society of Clinical Oncology

Creative Commons Attribution Non-Commercial No Derivatives 4.0 License 

INTRODUCTION:

Breast cancer (BC) is the most common type of cancer and the main cause of cancer-related death in women globally.¹ It has been estimated that there will be > 1.97 million new diagnoses of BC in women worldwide in 2020, and it is expected that 622,000 women will die of this disease.^{1,2}

Brazil is the largest country in Latin America and the fifth largest country in the world in terms of population.³ In Brazil, the prevalence and incidence of BC has progressively increased over the last years.⁴ BC is the main type of cancer among Brazilian women, according to the Instituto Nacional de Câncer. There were 59,700 new BC cases in 2018⁴ and there were

16,724 BC-associated deaths in 2017,⁵ with a higher mortality-to-incidence ratio when compared with developed countries.⁶ Previous work from our group demonstrated that Brazilian women have a higher risk of being diagnosed with late-stage BC and at a younger age than women in high-income countries.⁶⁻⁸ The 5-year overall survival (OS) rates previously reported for early-stage BC in Brazil were 96.84% for stage I and 94.16% for stage II disease. In locally advanced BC (stage III), the 5-year OS is 70.48%.⁷ Triple-negative and HER-2–positive BC subtypes were associated with inferior outcomes in terms of OS in patients with stage II or III BC.⁷

According to international guidelines, BC in young women (≤ 40 years) is an uncommon diagnosis, with

Author affiliations and support information (if applicable) appear at the end of this article.

Accepted on September 2, 2019 and published at ascopubs.org/journal/jgo on November 15, 2019; DOI <https://doi.org/10.1200/JGO.19.00263>

CONTEXT SUMMARY

Key Objective

Previous data indicate a higher prevalence of breast cancer (BC) in young women in Latin America compared with high-income countries. The purpose of this study was to characterize BC in young women in the Brazilian population through the AMAZONA III study.

Knowledge Generated

Of the women included in this representative prospective cohort, 17% were diagnosed with BC at ≤ 40 years of age. Young patients presented unfavorable clinicopathological features of BC at diagnosis, with more aggressive subtypes and advanced stage when compared with older women.

Relevance

We evaluated BC in young women at 22 Brazilian sites, including all regions of the country (public and private practice). The causes of a higher prevalence of BC among young women in Brazil deserves additional investigation since this could generate actions and public policies to reduce advanced-stage diagnosis of BC in this population.

a 0.40% to 0.45% cumulative risk of developing BC at the age of 40 years, which represents $< 7\%$ of all BC cases diagnosed in developed countries.⁹ In Latin American countries, BC is diagnosed at an earlier age when compared with patients from high-income countries, with a higher proportion of BC occurring among young women. Women younger than 40 years represent approximately 20% of the new cases of BC and 14% of the deaths resulting from BC.^{10,11}

In Brazil, the median age at BC diagnosis was reported to be 53 to 55 years⁶⁻⁸; however, the characteristics and prognosis of BC in young women in the Brazilian population are poorly described.¹²⁻¹⁵ Single-institution case series of young women with BC in Brazil have reported more advanced disease stages at the time of diagnosis, with a higher prevalence of metastatic disease.^{12,16,17}

AMAZONA III (ClinicalTrials.gov identifier: [NCT02669373](https://clinicaltrials.gov/ct2/show/study/NCT02669373)) is a prospective cohort study that includes patients with newly diagnosed BC from January 2016 to March 2018, involving 22 Brazilian sites and representing all five regions of the country. The main objective of this subanalysis was to characterize BC in young women in this population.

METHODS

This is a subanalysis of AMAZONA III study (GBECAM 0115),⁸ a prospective registry that included 2,950 women newly diagnosed with invasive BC in Brazil during the period of January 2016 to March 2018 within 22 sites, including patients covered by public and private health systems. Of these women, 2,888 had valid data regarding age at diagnosis and complete baseline information. To compare epidemiologic and clinicopathological features at the time of diagnosis of BC, patients were divided into two groups according to age: ≤ 40 years (group 1) and > 40 years (group 2). Ethnicity was defined according to the Brazilian Institute of

Geography and Statistics and race of the patients was classified as white, black, brown, indigenous, and Asian.¹⁸

BC subtypes were classified by immunohistochemistry for estrogen receptor (ER) status, progesterone receptor (PgR) status, and HER-2 status. All tumor tissue analysis and grading were performed in local laboratories. We used data from pathology reports to classify BC into five subtypes: luminal A (grade 1-2, ER and PgR positive, HER2 negative), luminal B (grade 3, ER and/or PgR positive, HER2 negative), luminal HER2 positive, nonluminal HER2 positive and triple negative (ER and PgR negative, HER2 negative). There was no central pathological review.

Quantitative variables were described as means, and categorical variables were described as frequencies and percentages and compared using the Pearson χ^2 test and adjusted residuals, when necessary. $P < .05$ was considered statistically significant. All analyses were performed using SAS, version 9.4 (SAS Institute, Cary, NC).

The AMAZONA III study was approved by the institutional review boards of each participating center and the national ethics committee.

RESULTS

Of 2,888 women diagnosed with invasive BC, we identified 486 (17%) who were ≤ 40 years. The baseline characteristics of younger and older patients are described and compared in [Tables 1 and 2](#).

Women aged ≤ 40 years were mostly of white ethnicity (54.2%), no statistical differences in health insurance coverage between the two age groups were found, and the majority of patients in both groups were insured by the public health system. The mean body mass index (calculated as kilograms divided by square meters) was 24.4 for patients ≤ 40 years and 24.9 for patients older than

40 years, with no statistical differences between the two groups. Also, there were no statistical differences in personal income, performance status, and family history of cancer between the two groups.

Women aged ≤ 40 years had significantly higher educational levels ($P < .001$), with group 1 versus group 2 data as follows: rates of illiteracy, 0.4% versus 5.6%; fewer than 8 years of school, 14.4% versus 30.4%; 9 to 11 years of

TABLE 1. Demographic and Socioeconomic Features by Age Groups of Brazilian Women Included in the AMAZONA III Study

Characteristic	Age at Diagnosis (years)		P
	≤ 40	> 40	
Race			
White	254 (54.2)	1,379 (58.7)	.051
Black	24 (5.1)	150 (6.4)	
Brown	183 (39.0)	803 (34.2)	
Indigenous	1 (0.2)	3 (0.1)	
Asian	7 (1.5)	14 (0.6)	
Health insurance			
Public	299 (62.0)	1,538 (64.7)	.270
Private	183 (38.0)	840 (35.3)	
Education			
Illiterate	2 (0.4)	121 (5.6)	$< .001$
< 8 years of school	64 (14.4)	663 (30.4)	
Completed 8 years of school	63 (14.2)	322 (14.8)	
Between 9 and 11 years of school	168 (37.7)	497 (22.8)	
University degree	148 (33.3)	574 (26.4)	
Personal income, US\$			
None	82 (23.8)	350 (19.9)	.236
Less than 1 minimum wage (< 233)	38 (11.0)	207 (11.7)	
1 to 2 minimum wages (880 to 467)	134 (38.8)	788 (44.7)	
2 to 3 minimum wages (467 to 701)	37 (10.7)	153 (8.7)	
3 to 5 minimum wages (701 to 1,167)	34 (9.9)	133 (7.5)	
5 to 10 minimum wages (1,167 to 2,338)	13 (3.8)	91 (5.2)	
10 to 20 minimum wages (2,338 to 4,670)	5 (1.4)	26 (1.5)	
> 20 minimum wages (> 4,670)	2 (0.6)	15 (0.9)	
Formal working activity			
Yes	272 (57.3)	924 (39.8)	$< .001$
No	203 (42.7)	1,398 (60.2)	
Currently married or lives in common-law marriage			
Yes	332 (69.2)	1,315 (56.5)	$< .001$
No	148 (30.8)	1,014 (43.5)	
Smoking history			
Never	352 (80.7)	1,448 (65.8)	$< .001$
Former	52 (11.9)	547 (24.9)	
Current	32 (7.3)	205 (9.3)	
Drinks alcoholic beverages			
Yes	127 (30.4)	479 (22.5)	$< .001$
No	291 (69.6)	1,648 (77.5)	
BMI, mean, kg/m ² (No.)	24.4 (486)	24.9 (2,400)	—

NOTE. Data presented as No. (%) unless otherwise indicated. Data reported in bold refers to $P < 0.05$.

Abbreviation: BMI, body mass index; US\$, US dollars.

TABLE 2. Reproductive Characteristics by Age Groups of Brazilian Women Included in the AMAZONA III Study

Characteristic	Age at Diagnosis		P
	≤ 40	> 40	
Mean age at menopause	43.5 (12)	47.8 (1,428)	NA
Mean age at menarche	12.6 (399)	13.1 (2,045)	NA
Reproductive status			
Premenopausal	339 (76.5)	415 (18.4)	< .001
Perimenopause	87 (19.6)	251 (11.1)	
Postmenopausal	17 (3.9)	1,588 (70.5)	
Ever used oral contraceptives			
Yes	346 (81.8)	1,314 (64.9)	< .001
No	77 (18.2)	711 (35.1)	
History of pregnancy			
Yes	383 (81.3)	1,973 (86.5)	.004
No	88 (18.7)	309 (13.5)	
Breastfeeding			
Yes	300 (87.5)	1,537 (87.0)	.809
No	43 (12.5)	230 (13.0)	
Diagnosed with breast cancer during a pregnancy			
Yes	12 (3.2)	9 (0.5)	< .001
No	364 (96.8)	1,881 (99.5)	

NOTE. Data presented as No. (%) unless otherwise indicated. Data reported in bold refers to $P < 0.05$. Abbreviation: NA, not applicable.

school, 37.7% versus 22.8%; and university degree, 33% versus 26.4%. Also, younger women (group 1) more frequently were employed than were older patients (57.3% v 39.8%; $P < .001$); and more frequently were married (69.2% v 56.5%; $P < .001$). In relation to social habits, women in group 1 smoked less (19.2% v 34.2%; $P < .001$) and consumed more alcohol than women in group 2 (30.4% v 22.5%; $P < .001$).

There were significant differences regarding previous use of contraceptives in (81.8% v 35.1%; $P < .001$) and nulliparity (18.7% v 13.5%; $P = 0.004$) in younger and older patients. Twelve women ≤ 40 years (3.2%) had a diagnosis of BC during pregnancy, compared with nine patients (0.5%) in the older group ($P < .001$).

The mode how BC was detected differed significantly between the two groups: BC in younger women was more often detected symptomatically than in older women (73.4% v 64.5%; $P < .001$; Table 3). Initial tumor size was also significantly larger among younger women ($P < .001$). The prevalence of T1, T2, T3, and T4 tumors at the time of diagnosis for women in group 1 compared with those in group 2 were, respectively: 27.1% versus 36.9%, 33.6% versus 37.6%, 24.1% versus 13.9%, and 15.2% versus 11.6% (Fig 1C; Table 3). There was no statically significant difference in initial positive lymph node status between the two groups (44.1% and 39%, for group 1 and group 2, respectively; $P = .152$; Data Supplement).

Stage I disease at diagnosis occurred in 19.2% of patients aged ≤ 40 years and 27.8% in women > 40 years old ($P < .001$). In contrast, stage III disease was found in 36.8% of younger women and 25.1% of older women ($P < .001$; Fig 1A; Data Supplement). Women in group 1 had more grade 3 tumors (43.1% v. 30.1%; $P < .001$; Fig 1B; Data Supplement). Younger women more frequently underwent mastectomy than did older women (54.7% v 45.7%, $P < .001$; Data Supplement), and this difference remained statistically significant when adjusting for initial staging (Mantel-Haenszel method).

The distribution of BC subtypes, as determined by immunohistochemistry, was distinct among young and older women (Fig 2; Data Supplement). Younger patients had a higher proportion of luminal B HER2-negative (15.8% v 11.4%; $P < .001$), luminal B HER2-positive (22.8% v 16%; $P < .001$), and triple-negative (23% v 14.1%; $P < .001$) tumors at diagnosis (Fig 2; Data Supplement).

DISCUSSION

BC incidence is growing in Latin America and it is considered an important health burden.^{10,11,19} There is limited information about the diagnosis, treatment patterns, and outcomes of young women with BC in Latin American countries. To our knowledge, AMAZONA III is the first prospective, multicentric registry of BC in Brazil; its objective is to better describe the current scenario of BC care in the largest and most populated country in Latin

TABLE 3. Patients at Baseline: Tumor Characteristics and Type of Surgery by Age Groups of Brazilian Women Included in the AMAZONA III Study

Characteristic	Age at Diagnosis (years)		P
	≤ 40	> 40	
ECOG performance stage			
0	285 (81.0)	1,358 (74.7)	.080
1	63 (17.9)	405 (22.3)	
2	3 (0.8)	45 (2.4)	
3	1 (0.3)	7 (0.4)	
4	0 (0.0)	3 (0.2)	
Diagnostic			
Screening-detected	120 (26.6)	807 (35.5)	.003
Symptomatic	331 (73.4)	1,466 (64.5)	
TNM			
T1	114 (27.1)	749 (36.9)	< .001
T2	141 (33.6)	764 (37.6)	
T3	101 (24.1)	282 (13.9)	
T4	64 (15.2)	235 (11.6)	
Stage at diagnosis			
I	76 (19.2)	541 (27.8)	< .001
II	156 (39.4)	816 (41.9)	
III	146 (36.8)	489 (25.1)	
IV	18 (4.6)	101 (5.2)	
Histology			
Ductal	392 (94.7)	1,793 (87.6)	< .001
Lobular	15 (3.6)	171 (8.3)	
Mucinous	0 (0.0)	30 (1.5)	
Papillary	2 (0.5)	20 (1.0)	
Medullary	3 (0.7)	4 (0.2)	
Mixed	2 (0.5)	29 (1.4)	
KI-67, mean (No.)	44.7 (355)	30.3 (1,984)	—
Tumor grade			
1	46 (10.7)	381 (17.9)	< .001
2	198 (46.2)	1,110 (52.0)	
3	185 (43.1)	641 (30.1)	
HER-2 status			
Positive	107 (29.6)	438 (22.3)	.003
Negative	255 (70.4)	1,526 (77.7)	
Hormone receptor			
Positive	273 (71.1)	1,646 (79.4)	.001
Negative	111 (28.9)	428 (20.6)	

(Continued in next column)

TABLE 3. Patients at Baseline: Tumor Characteristics and Type of Surgery by Age Groups of Brazilian Women Included in the AMAZONA III Study (Continued)

Characteristic	Age at Diagnosis (years)		P
	≤ 40	> 40	
Breast cancer subtype			
Luminal A	106 (30.6)	957 (51.3)	< .001
Luminal B, HER-2 negative	55 (15.8)	212 (11.4)	
Luminal B, HER-2 positive	79 (22.8)	298 (16.0)	
HER-2 positive	27 (7.8)	135 (7.2)	
Triple negative	80 (23.0)	264 (14.1)	
Lymph node			
Positive	94 (44.1)	541 (39.0)	.152
Negative	119 (55.9)	847 (61.0)	
Surgery			
Breast-conserving surgery	82 (38.7)	708 (51.3)	
Mastectomy	116 (54.7)	630 (45.7)	< .001
Adenomastectomy	8 (3.8)	21 (1.5)	
Skin-sparing mastectomy	6 (2.8)	20 (1.5)	

NOTE. Data presented as No. (%) unless otherwise indicated. Data reported in bold refers to $P < 0.05$.

Abbreviation: ECOG, Eastern Cooperative Oncology Group.

America. The increased proportion of BC among young women is of utmost importance, since they tend to be diagnosed in later stages and usually bear more aggressive BC subtypes that, at long run, can negatively impact their survival.^{9,11,20}

We have shown that young women diagnosed with BC are more frequently employed and actively working, have a higher educational level, and most are married, highlighting the socioeconomic impact BC may have when afflicting young women. It is well known that BC and its treatment can largely affect the economy of a community, because some BC survivors experience reduced work ability.²¹⁻²³ Recently, Landeiro et al²⁴ reported return to work (RTW) rates of 30.3% and 60.4% at 12 and 24 months, respectively, after BC diagnosis in a cohort of 125 employed women from a single institution in Brazil. These rates are lower when compared with affected women in high-income countries. Because the cohort of young women in the AMAZONA III study represents an important productive working force from several regions of the country, additional follow-up of the RTW rates and the factors associated with the RTW decision in this population is paramount.

BC in young women is considered a somewhat rare situation, accounting for 5% to 7% of BC cases in developed countries.^{9,20,25,26} In our cohort, women aged ≤ 40 years represented 17% of patients, a number significantly higher than the rates reported in the literature. In our previous work, we have reported that the median age at




FIG 1. Bar graphs of (A) clinical stage, (B) tumor grade, and (C) tumor size at diagnosis by age groups of Brazilian women included in the AMAZONA III study. (*) $P < .001$.

diagnosis of BC in Brazil is 53 to 55 years,⁶⁻⁸ whereas in high-income countries it is approximately 64 years.^{1,27} Also, a review of the literature undertaken by Villarreal-Garza et al¹¹ reported a higher BC incidence and mortality rate in young women in Latin America when compared with those in developed countries (20% v. 12% and 14% v. 7%, respectively).

Another study, led by Franco-Marina et al,¹⁹ confirmed the high incidence of BC in young women in Latin America, estimating that one in every five cases of BC is diagnosed in women younger than 45 years, almost double of the frequency observed in developed countries like the United

States and Canada. This high incidence of cancer in young women remained statistically significant even after adjusting for the percentage of young women in Latin and North American populations. All these studies suggest a high prevalence of young patients with BC in Latin America, and the causes should be better investigated.

Regarding the potential modifiable risk factors for BC to develop in young women, we recognized in our cohort a significantly higher prevalence of alcohol consumption among young women. Recently, an effort has been made by public health experts and medical societies to emphasize that reducing alcohol consumption is a vital, and largely




FIG 2. Breast cancer subtype by age groups of Brazilian women included in the AMAZONA III study. (*) $P < .001$.

neglected, cancer prevention strategy.²⁸⁻³⁰ Greater attention to this matter should be given to effectively communicate the role of alcohol as a risk factor for BC and to better investigate its impact on the prevalence of BC in Latin America.

Women younger than 40 years are usually diagnosed with more aggressive disease and have worse prognosis (based on more advanced clinical stage and worse histopathological characteristics, with a high prevalence of high-grade tumors), and they have a higher risk of recurrence when compared with older women.^{11,16,19,31} This is especially observed in luminal BC subtypes, for which young women have inferior outcomes compared with older women with the same initial tumor characteristics.³² For these reasons, younger women sometimes receive more aggressive treatment than do older women.^{17,20,33}

Histopathological and immunohistochemical characteristics presented in our study also corroborate the data from the literature,^{11,17,33,34} with a larger number of locally advanced lesions (T3, T4) and poorly differentiated tumors among young women. Although the luminal subtype is still the most frequent among young women, we reported significant differences in the frequency of the luminal subtype between older and younger women, with more luminal B type (15.8% v 11.4%, group 1 v group 2, respectively) or luminal B/HER-2 positive (22.8% v 16%, group 1 v group 2, respectively) in the latter group. In comparison with series of young women from the United States^{35,36} and Europe,³⁷ our population had a greater proportion of triple-negative subtype BC (23% v 14.1%). In a previous retrospective, single-center cohort report (N = 738), a lower proportion of ER-positive tumors were reported in younger women as compared with older women (33.5% v 42.8%), and higher proportion of triple-negative BC (10% v 6.4%) also was reported among young women (n = 376) as compared with older women.³⁴

In our study, we observed striking differences regarding stage distribution between the two age groups. Stage I disease was more frequent among older women than younger women (27.8% v 19.2%, respectively), whereas younger women had significantly more stage III disease (36.9%) than older women (25.1%). Metastatic BC de novo was present in 4.5% of the women \leq 40 years old and 5.2% of older women. Indeed, 41.4% of the young women in our study were initially diagnosed with stage III or IV BC. Our data are comparable to that of another large cohort of Latin American women, from Mexico, in which 15% of women with BC were \leq 40 years old.³³ Also, young Mexican women had more aggressive disease at presentation, with tumors of higher grade and a larger proportion of luminal B and triple-negative cancers.³³ Similar to our data, fewer stage I tumors were found in young women and more stage III and IV disease than in their older counterparts.³³

Another important issue to highlight is that in Brazil, there is a gap regarding the access to optimal therapy between patients in the public and private health systems, and this influences patient outcomes.^{38,39} This disparity will also play an important role in young women diagnosed with advanced disease (stage III and IV), because access to optimal HER2 therapy, ovarian suppression, and CDK 4/6 inhibitors might be restricted to patients in the private health system.

Finally, BC was detected by screening in only 26% of patients \leq 40 years old, as compared with 35.5% of older women. This finding is not unexpected, because the current national BC screening guideline recommends mammography starting at age 50 years.⁴⁰ However, they call attention to the low proportion of older women whose diagnosis is established through screening programs in Brazil. Better educational strategies, access to and compliance with screening programs are urgently needed in

Brazil. Organized screening of young, healthy women has been controversial, regarded as inefficient and even deleterious by some experts, and has not been recommended widely.⁴¹ Individualized screening, targeting only some high-risk young women, may be beneficial, although no randomized trial has shown an impact on BC mortality.⁴² All recommendations for this younger age group are based on experts' opinions and do not take into account the higher proportion of BC in young women found in our study and in other Latin American cohorts. In these populations, earlier screening should be a subject of additional research.

This study was of a large cohort of young women with BC in Brazil and was a prospective evaluation including patients from all regions of the country, from the private and the public health systems. Longer follow-up of this population will reveal important information regarding treatment patterns and outcomes. Our study has some limitations, especially regarding BC subtypes, because we did not perform central revision of the immunohistochemistry data.

At this time, we are not aware of any specific clinical or educational programs focused on this population in Brazil. Particular aspects of the care for young women with BC have been studied and reported in the literature. It would be interesting if at least the major reference centers of cancer

care in Brazil could implement specific units for the care of young women with BC, addressing the particular needs of this growing population (eg, fertility preservation, precise psychosocial interventions, genetic evaluation, aspects related to corporal image and reconstructive surgery support, challenges related to a longer survival and follow-up period). Successful examples of this concept include centers in the United States (Young and Strong Program for Young Women with Breast Cancer),⁴³ Canada (Breast Cancer Program for Young Women),⁴⁴ and Mexico (Joven and Fuerte: Program for Young Women with Breast Cancer in Mexico).⁴⁵ A call to action from health policy planners, medical providers, researchers, patients with BC, their families, and the community in general is encouraged for better care of this emergent challenge.

In conclusion, in Brazil, a higher proportion of patients with BC are diagnosed at ≤ 40 years of age when compared with women in developed countries. Younger patients have unfavorable clinicopathological features, with advanced stages and more aggressive BC subtypes at diagnosis, when compared with older patients. The causes of a high prevalence of BC in young women should be investigated further, as should potential preventive and screening strategies.

AFFILIATIONS

¹Hospital de Clínicas de Porto Alegre, Porto Alegre, Brazil

²Jules Bordet Institut, Brussels, Belgium

³Latin American Cooperative Oncology Group, Porto Alegre, Brazil

⁴Grupo Brasileiro de Estudos em Câncer de Mama, Porto Alegre, Brazil

⁵Hospital Moinhos de Vento, Porto Alegre, Brazil

⁶Instituto Nacional do Câncer, Rio de Janeiro, Brazil

⁷Centro Regional Integrado de Oncologia, Fortaleza, Brazil

⁸Hospital Araújo Jorge, Goiás, Brazil

⁹A.C. Camargo Cancer Center, São Paulo, Brazil

¹⁰Universidade Federal de Goiás, Goiânia, Brazil

¹¹Hospital do Câncer de Londrina, Londrina, Brazil

¹²Liga Norte Riograndense, Natal, Brazil

¹³Hospital Jardim Amália, Volta Redonta, Brazil

¹⁴Oncoclínica Rio de Janeiro, Rio de Janeiro, Brazil

¹⁵Centro de Pesquisa e Educação da Serra Gaúcha- DeVita, Caxias do Sul, Brazil

¹⁶Instituto do Câncer Dr Arnaldo, São Paulo, Brazil

¹⁷Centro de Pesquisas em Oncologia, Florianópolis, Brazil

¹⁸Clínica Assistência Multidisciplinar em Oncologia, Salvador, Brazil

¹⁹Centro Integrado de Terapia Onco-Hematológica, Passo Fundo, Brazil

²⁰Universidade de Caxias do Sul, Caxias do Sul, Brazil

²¹Instituto Oncológico de Ribeirão Preto, Ribeirão Preto, Brazil

²²Núcleo Oncológico da Bahia, Lauro de Freitas, Brazil

²³Hospital São Vicente de Paulo, Passo Fundo, Brazil

²⁴Clínica de Neoplasia Litoral, Itajaí, Brazil

²⁵Instituto do Câncer do Estado de São Paulo, São Paulo, Brazil

CORRESPONDING AUTHOR

Maria Alice Franzoi, MD, Clinical Trials Research Unit, Institut Jules Bordet, Boulevard Waterloo, 121, Brussels, Belgium, 1000; e-mail: franzoi.alice@gmail.com.

SUPPORT

Supported by the Latin American Cooperative Oncology Group (M.A.F.) and Grupo Brasileiro de Estudos em Câncer de Mama (M.A.F.).

AUTHOR CONTRIBUTIONS

Conception and design: Maria Alice Franzoi, Facundo Zaffaroni, Gustavo Werutsky, Carlos Barrios, Ruffo Freitas Júnior, Yeni Nerón, Clarissa Mathias, Giuliano Borges, Max Mano, Pedro E. R. Liedke

Financial support: Yeni Nerón

Administrative support: Yeni Nerón

Provision of study material or patients: Eduardo Cronemberger, Geraldo Silva Queiroz, Vladimir Cordeiro de Lima, Ruffo Freitas Júnior, José Couto, Heloísa Resende, Susana Crocamo, Tomás Reinert, Yeni Nerón, Vanessa Dybal, Rita de Cassia Costamillan, Alessandra Morelle, Carlos Alberto Sampaio Filho

Collection and assembly of data: Maria Alice Franzoi, Facundo Zaffaroni, Gustavo Werutsky, José Bines, Carlos Barrios, Eduardo Cronemberger, Geraldo Silva Queiroz, Vladimir Cordeiro de Lima, Ruffo Freitas Júnior, Karla Emerenciano, Heloísa Resende, Susana Crocamo, Tomás Reinert, Brigitte Van Eyli, Yeni Nerón, Vanessa Dybal, Nicolas Lazaretti, Rita de Cassia Costamillan, Diocésio Alves Pinto de Andrade, Clarissa Mathias, Giovana Zerwes Vacaro, Alessandra Morelle, Carlos Alberto Sampaio Filho

Data analysis and interpretation: Maria Alice Franzoi, Daniela D. Rosa, Facundo Zaffaroni, Gustavo Werutsky, Sérgio Simon, Vladimir Cordeiro de Lima, Ruffo Freitas Júnior, José Couto, Yeni Nerón, Clarissa Mathias, Alessandra Morelle, Max Mano, Pedro E. R. Liedke

Manuscript writing: All authors

Final approval of manuscript: All authors

Accountable for all aspects of the work: All authors

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated unless otherwise noted. Relationships are self-held unless noted. I = Immediate

Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to www.asco.org/rwc or ascopubs.org/jgo/site/misc/authors.html.

Open Payments is a public database containing information reported by companies about payments made to US-licensed physicians ([Open Payments](http://OpenPayments.gov)).

Daniela D. Rosa

Honoraria: Roche, Novartis, Pfizer, AstraZeneca, Eisai, Eli Lilly

Consulting or Advisory Role: Eli Lilly, Novartis, Roche

Sérgio Simon

Consulting or Advisory Role: Roche, AstraZeneca, MSD Oncology, Pfizer

Speakers' Bureau: AstraZeneca, Novartis, Roche

Travel, Accommodations, Expenses: Roche, AstraZeneca

Carlos Barrios

Stock and Other Ownership Interests: Biomarker, MedSIR, Tummi

Honoraria: Novartis, Roche, Pfizer, GlaxoSmithKline, Sanofi, Boehringer Ingelheim, Eisai

Consulting or Advisory Role: Boehringer Ingelheim, Roche, Novartis, GlaxoSmithKline, Eisai, Pfizer, AstraZeneca, Libbs, MSD Oncology, United Medical

Research Funding: Pfizer, Novartis, Amgen, AstraZeneca, Boehringer Ingelheim, GlaxoSmithKline, Roche, Eli Lilly, Sanofi, GlaxoSmithKline, Taiho Pharmaceutical, Mylan, Merrimack, Merck, AbbVie, Astellas Pharma, Biomarin, Bristol-Myers Squibb, Daiichi Sankyo, Abraxis BioScience, AB Science, Asana Biosciences, Medivation, Daiichi Sankyo, Exelixis, ImClone Systems, LEO Pharma, Millennium, Janssen, Atlantis Clinica, INC Research, Halozyme, Covance, Celgene, inVentiv Health

Travel, Accommodations, Expenses: Roche, Novartis, Pfizer, BMS Brazil, AstraZeneca, MSD Oncology

Eduardo Cronemberger

Honoraria: Roche, Merck Sharp & Dohme

Consulting or Advisory Role: Roche, Merck Sharp & Dohme

Research Funding: AstraZeneca/MedImmune (Inst), Roche (Inst), MSD Oncology (Inst), Janssen-Cilag (Inst), Merck Serono (Inst)

Travel, Accommodations, Expenses: AstraZeneca, Pfizer

Vladimir Cordeiro de Lima

Honoraria: Roche, BMS Brazil, MSD Oncology, AstraZeneca, Boehringer Ingelheim, Pfizer, Merck Serono

Consulting or Advisory Role: Roche, BMS Brazil, AstraZeneca, Pfizer, MSD Oncology

Speakers' Bureau: Roche, BMS Brazil, Boehringer Ingelheim, MSD Oncology, Merck Serono

Travel, Accommodations, Expenses: Roche, MSD Oncology, AstraZeneca, Boehringer Ingelheim, BMS Brazil

Ruffo Freitas Júnior

Honoraria: Roche, Libbs, Pfizer

Consulting or Advisory Role: Roche

Research Funding: Roche (Inst), MSD Oncology (Inst)

Travel, Accommodations, Expenses: Roche, Libbs

Heloísa Resende

Travel, Accommodations, Expenses: Pfizer

Tomás Reinert

Honoraria: Novartis, AstraZeneca, Pfizer, Eli Lilly, Pierre Fabre, AstraZeneca

Yeni Nerón

Other Relationship: AstraZeneca (Inst), Roche (Inst), Clovis Oncology (Inst), Merck Serono (Inst)

Vanessa Dybal

Speakers' Bureau: Novartis, Roche

Travel, Accommodations, Expenses: Pfizer, Roche

Nicolas Lazaretti

Research Funding: Acerta Pharma (Inst), BeiGene (Inst), Bristol-Myers Squibb (Inst), AstraZeneca (Inst), Roche (Inst), Merck (Inst), Medivation (Inst), Regeneron (Inst)

Travel, Accommodations, Expenses: MSD Oncology, Roche

Diocésio Alves Pinto de Andrade

Speakers' Bureau: Novartis

Clarissa Mathias

Consulting or Advisory Role: Roche

Alessandra Morelle

Consulting or Advisory Role: Roche, Amgen, MSD Oncology, Eli Lilly, Novartis

Speakers' Bureau: Roche, Eli Lilly, Novartis, MSD Oncology

Carlos Alberto Sampaio Filho

Employment: Clinica AMO

Leadership: Clinica AMO Salvador Brazil

Stock and Other Ownership Interests: Clinica AMO

Consulting or Advisory Role: Pneuma Respiratory

Max Mano

Stock and Other Ownership Interests: Biotoscana, Fleury Group, Hypera

Honoraria: Roche, Novartis, Oncologia Brasil, Eli Lilly/ImClone

Consulting or Advisory Role: Novartis, Roche, Pfizer, AstraZeneca, Eli Lilly/ImClone

Travel, Accommodations, Expenses: Roche

Uncompensated Relationships: Roche

Pedro E. R. Liedke

Consulting or Advisory Role: Zodiac Pharma

Research Funding: Merck Sharp & Dohme, Merck Serono, Roche, AstraZeneca, Halozyme, Acerta Pharma, Novartis, Bristol-Myers Squibb, Regeneron, Medivation, Myovant Sciences, Janssen, PharmaMar, Eurofarma, Pfizer, PPD, PRA Health Sciences, Covance, Quintiles, ICON Clinical Research, Parexel, Intrials

Travel, Accommodations, Expenses: Pfizer, Roche, Oncoclinicas

No other potential conflicts of interest were reported.

ACKNOWLEDGMENT

Latin America Cooperative Oncology Group thanks Itaú, Avon, SAS Data4Good, and Projeto CURA for their support.

REFERENCES

1. Bray F, Ferlay J, Soerjomataram I, et al: Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 68:394-424, 2018
2. Siegel RL, Miller KD, Jemal A: Cancer statistics, 2018. *CA Cancer J Clin* 68:7-30, 2018
3. World Bank: DataBank. World development indicators. <https://databank.worldbank.org/reports.aspx?source=2&series=SP.POP.TOTL&country=WLD>

4. Instituto Nacional de Câncer José Alencar Gomes da Silva. Coordenação de Prevenção e Vigilância: Estimativa 2018: Incidência de Câncer no Brasil. <https://www.inca.gov.br/publicacoes/livros/estimativa-2018-incidencia-de-cancer-no-brasil>
5. Instituto Nacional de Câncer, Ministério da Saúde: Atlas On-line de Mortalidade. <https://mortalidade.inca.gov.br/MortalidadeWeb/>
6. Lee BL, Liedke PE, Barrios CH, et al: Breast cancer in Brazil: Present status and future goals. *Lancet Oncol* 13:e95-e102, 2012
7. Simon SD, Bines J, Werutsky G, et al: Characteristics and prognosis of stage I-III breast cancer subtypes in Brazil: The AMAZONA retrospective cohort study. *Breast* 44:113-119, 2019
8. Rosa D, Barrios C, Bines J, et al: Abstract P1-08-29: Current status of clinical and pathological characteristics of breast cancer patients in Brazil: Results of the AMAZONA III study (GBECAM 0115). *Cancer Res* 79, 2019 (abstr P1-08-29)
9. Paluch-Shimon S, Pagani O, Partridge AH, et al: ESO-ESMO 3rd International Consensus Guidelines for Breast Cancer in Young Women (BCY3). *Breast* 35:203-217, 2017
10. Cazap E: Breast cancer in Latin America: A map of the disease in the region. *Am Soc Clin Oncol Educ Book* 38:451-456, 2018
11. Villarreal-Garza C, Aguila C, Magallanes-Hoyos MC, et al: Breast cancer in young women in Latin America: An unmet, growing burden. *Oncologist* 18:1298-1306, 2013
12. Rocha-Brischiliari SC, Oliveira RR, Andrade L, et al: The rise in mortality from breast cancer in young women: Trend analysis in Brazil. *PLoS One* 12:e0168950, 2017
13. Cecilio AP, Takakura ET, Jumes JJ, et al: Breast cancer in Brazil: Epidemiology and treatment challenges. *Breast Cancer (Dove Med Press)* 7:43-49, 2015
14. Fujimoto RHP, Koifman RJ, Silva IFD: Survival rates of breast cancer and predictive factors: A hospital-based study from western Amazon area in Brazil. *Cien Saude Colet* 24:261-273, 2019
15. Di Sibio A, Abriata G, Forman D, et al: Female breast cancer in Central and South America. *Cancer Epidemiol* 44:S110-S120, 2016 (Suppl 1)
16. Eugênio DSG, Souza JA, Chojniak R, et al: Breast cancer features in women under the age of 40 years. *Rev Assoc Med Bras (1992)* 62:755-761, 2016
17. Costa NCD, Morsch DM, Opperman CP, et al: Biological features of breast cancer according to age at diagnosis in southern Brazil: An analysis of retrospective data of 1128 women. *Breast J* 25:760-762, 2019
18. Instituto Brasileiro de Geografia e Estatística: https://ww2.ibge.gov.br/english/estatistica/populacao/caracteristicas_raceis/default_raceis.shtm
19. Franco-Marina F, López-Carrillo L, Keating NL, et al: Breast cancer age at diagnosis patterns in four Latin American Populations: A comparison with North American countries. *Cancer Epidemiol* 39:831-837, 2015
20. Azim HA, Jr, Partridge AH: Biology of breast cancer in young women. *Breast Cancer Res* 16:427, 2014
21. Arfi A, Baffert S, Soilly A-L, et al: Determinants of return at work of breast cancer patients: Results from the OPTISOINS01 French prospective study. *BMJ Open* 8:e020276, 2018
22. Grinshpun A, Rottenberg Y: Unemployment following breast cancer diagnosis: A population-based study. *Breast* 44:24-28, 2019
23. Heuser C, Halbach S, Kowalski C, et al: Sociodemographic and disease-related determinants of return to work among women with breast cancer: A German longitudinal cohort study. *BMC Health Serv Res* 18:1000, 2018
24. Landeiro LCG, Gagliato DM, Fêde AB, et al: Return to work after breast cancer diagnosis: An observational prospective study in Brazil. *Cancer* 124:4700-4710, 2018
25. Assi HA, Khoury KE, Dbouk H, et al: Epidemiology and prognosis of breast cancer in young women. *J Thorac Dis* 5(Suppl 1):S2-S8, 2013
26. Anastasiadi Z, Lianos GD, Ignatiadou E, et al: Breast cancer in young women: An overview. *Updates Surg* 69:313-317, 2017
27. DeSantis C, Ma J, Bryan L, et al: Breast cancer statistics, 2013. *CA Cancer J Clin* 64:52-62, 2014
28. Chambers SE, Copson ER, Dutey-Magni PF, et al: Alcohol use and breast cancer risk: A qualitative study of women's perspectives to inform the development of a preventative intervention in breast clinics. *Eur J Cancer Care (Engl)* 28:e13075, 2019
29. LoConte NK, Brewster AM, Kaur JS, et al: Alcohol and cancer: A statement of the American Society of Clinical Oncology. *J Clin Oncol* 36:83-93, 2018
30. Sinclair J, McCann M, Sheldon E, et al: The acceptability of addressing alcohol consumption as a modifiable risk factor for breast cancer: A mixed method study within breast screening services and symptomatic breast clinics. *BMJ Open* 9:e027371, 2019
31. Garicochea B, Morelle A, Andrighetti AE, et al: Idade como fator prognóstico no câncer de mama em estágio inicial [in Portuguese]. *Rev Saude Publica* 43:311-317, 2009
32. Partridge AH, Hughes ME, Warner ET, et al: Subtype-dependent relationship between young age at diagnosis and breast cancer survival. *J Clin Oncol* 34:3308-3314, 2016
33. Villarreal-Garza C, Mohar A, Bargallo-Rocha JE, et al: Molecular subtypes and prognosis in young Mexican women with breast cancer. *Clin Breast Cancer* 17:e95-e102, 2017
34. De Lima Vazquez F, Silva TB, Da Costa Vieira RA, et al: Retrospective analysis of breast cancer prognosis among young and older women in a Brazilian cohort of 738 patients, 1985-2002. *Oncol Lett* 12:4911-4924, 2016
35. Keegan THM, Press DJ, Tao L, et al: Impact of breast cancer subtypes on 3-year survival among adolescent and young adult women. *Breast Cancer Res* 15:R95, 2013
36. Bharat A, Aft RL, Gao F, et al: Patient and tumor characteristics associated with increased mortality in young women (< or =40 years) with breast cancer. *J Surg Oncol* 100:248-251, 2009
37. Cancellato G, Maisonneuve P, Rotmensz N, et al: Prognosis and adjuvant treatment effects in selected breast cancer subtypes of very young women (<35 years) with operable breast cancer. *Ann Oncol* 21:1974-1981, 2010
38. Debiasi M, Reinert T, Kaliks R, et al: Estimation of premature deaths from lack of access to anti-HER2 therapy for advanced breast cancer in the Brazilian public health system. *J Glob Oncol* 3:201-207, 2016
39. Liedke PER, Finkelstein DM, Szymonifka J, et al: Outcomes of breast cancer in Brazil related to health care coverage: A retrospective cohort study. *Cancer Epidemiol Biomarkers Prev* 23:126-133, 2014
40. Migowski A, Silva GAE, Dias MBK, et al: Diretrizes para detecção precoce do câncer de mama no Brasil. II - Novas recomendações nacionais, principais evidências e controvérsias [in Portuguese]. *Cad Saude Publica* 34:e00074817, 2018
41. Siu AL: US Preventive Services Task Force: Screening for breast cancer: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med* 164:279-296, 2016
42. Desreux JAC: Breast cancer screening in young women. *Eur J Obstet Gynecol Reprod Biol* 230:208-211, 2018
43. Partridge AH, Ruddy KJ, Kennedy J, et al: Model program to improve care for a unique cancer population: Young women with breast cancer. *J Oncol Pract* 8:e105-e110, 2012
44. Ali A, Warner E: pynk: Breast cancer program for young women. *Curr Oncol* 20:e34-e39, 2013
45. Villarreal-Garza C, Castro-Sánchez A, Platas A, et al: "Joven & Fuerte": Program for young women with breast cancer in Mexico - initial results. *Rev Invest Clin* 69:223-228, 2017

